

Effect of the unpolarized spin state in spin-correlation measurement of two protons produced in the $^{12}\text{C}(d,^2\text{He})$ reaction

February 1, 2008

S. Hamieh

*Kernfysisch Versneller Instituut,
Zernikelaan 25, 9747 AA Groningen, The Netherlands.*

Abstract

In this note we discuss the effect of the unpolarized state in the spin-correlation measurement of the $^1S_0$ two-proton state produced in $^{12}\text{C}(d,^2\text{He})$ reaction at the KVI, Groningen. We show that in the presence of the unpolarized state the *maximal* violation of the CHSH-Bell inequality is lower than the classical limit if the purity of the state is less than $\sim 70\%$. In particular, for the KVI experiment the violation of the CHSH-Bell inequality should be corrected by a factor $\sim 10\%$ from the pure $^1S_0$ state.

1 Introduction

In an experiment performed at the Kernfysisch Versneller Instituut (KVI), Groningen [5] with the goal to test Bell inequality violation in Nuclear Physics (perhaps to be applied in quantum information physics), the experimental group, by bombarding a ^{12}C target with 170 MeV d , was able to generate a singlet-spin, two-proton state coupled to unpolarized state with $\sim 10\%$ contribution. In this paper we will analyze the experimental results of this experiment and we will show that the effect of the unpolarized state is important and could not be neglected.

2 CHSH inequalities and entanglement in a mixed ensemble

Bell-type inequalities relating averages of four random dichotomic variables a , a' and b , b' representing measurements of spin in directions $\hat{a}$, $\hat{a}'$ and $\hat{b}$, $\hat{b}'$. The Clauser, Horne, Shimony and Holth (CHSH) [3] form of Bell-type inequalities for spin 1/2 case could be written in this form

$$|E(\phi_1, \phi'_1, \phi_2, \phi'_2)| = |E(\phi_1, \phi_2) + E(\phi_1, \phi'_2) + E(\phi'_1, \phi_2) - E(\phi'_1, \phi'_2)| \leq 2, \quad (1)$$

where ϕ_i is the analyzer angular setting for the i^{th} particles ($i = 1, 2$) and $E(\phi_i, \phi_j)$ is the correlation function defined as

$$E(\phi_i, \phi_j) = \frac{N_{++} + N_{--} - N_{+-} - N_{-+}}{N_{total}}. \quad (2)$$

In quantum-theory language the CHSH operator corresponding to the CHSH inequality is represented by an operator

$$\mathcal{B} = \hat{a} \cdot \sigma \otimes (\hat{b} + \hat{b}') \cdot \sigma + \hat{a}' \cdot \sigma \otimes (\hat{b} - \hat{b}') \cdot \sigma, \quad (3)$$

acting in Hilbert space $\mathcal{H}_A \otimes \mathcal{H}_B$ in $2 \otimes 2$ dimension. The correlation function is given by the mean value of the operator $\hat{a} \cdot \sigma \otimes \hat{b} \cdot \sigma$. For a pure state this correlation function could be easily computed, *e.g.* for singlet state we have

$$E(\phi_i, \phi_j) = -\cos(\phi_i - \phi_j). \quad (4)$$

For mixed state, however, the mean value should be averaged over the ensemble and therefore the CHSH inequality not a sufficient condition to test the presence of entanglement[7]. Different measures of the entanglement have been proposed in the literature for mixed state¹, *e.g.* entanglement of formation, distillation, relative entropy of entanglement, negativity, etc. . . Here we will use the entanglement of formation as our measure of the entanglement.

¹Any measurement of the entanglement should not increase by local operation (*e.g.* unitary transformation) and classical communication (*e.g.* phone calls.), known as LOCC

In a mixed ensemble any bipartite quantum state ρ_{AB} can be written as:

$$\rho_{AB} = \frac{1}{4} \left(I \otimes I + \mathbf{A} \cdot \boldsymbol{\sigma} \otimes I + I \otimes \mathbf{P} \cdot \boldsymbol{\sigma} + \sum_{i,j=1}^3 D_{ij} \sigma_i \otimes \sigma_j \right). \quad (5)$$

σ_i are the pauli matrices, I is the identity operator, $\mathbf{A}$ and $\mathbf{P}$ are vectors in $\mathcal{R}^3$. The D_{ij} form a 3×3 matrix called the correlation matrix D . In this representation of the density matrix the mean value of the CHSH-Bell operator is given by [4]

$$\langle \mathcal{B} \rangle = \hat{a} \cdot [D(\hat{b} + \hat{b}')] + \hat{a}' \cdot [D(\hat{b} - \hat{b}')] . \quad (6)$$

Using the representation of the density matrix given in Eq. (5), we characterize any bipartite quantum state ρ_{AB} by

- The entanglement measured by the “tangle”, τ , of the entanglement of formation [6] and defined by

$$\tau = \max\{\lambda_1 - \lambda_2 - \lambda_3 - \lambda_4, 0\}, \quad (7)$$

where the λ 's are the square roots of the eigenvalues, in decreasing order, of the matrix, $\rho_{AB}(\sigma_y \otimes \sigma_y \rho_{AB}^* \sigma_y \otimes \sigma_y)$ and ρ_{AB}^* is the complex conjugation of ρ_{AB} in the computational basis $\{|++\rangle, |+-\rangle, |-+\rangle, |--\rangle\}$.

- The maximum amount of the CHSH-Bell violation of the state ρ_{AB} [4]

$$\langle \mathcal{B} \rangle^{max} = 2\sqrt{M(\rho_{AB})}. \quad (8)$$

$M(\rho_{AB})$ is the sum of the two larger eigenvalues of $DD^\dagger$ ².

- The purity of the state that measures how far the state is from pure state

$$S_L = \text{Tr}(\rho_{AB}^2). \quad (9)$$

3 Analysis of the experimental data of the KVI experiment

The spin state of the two protons produced in the $^{12}\text{C}(d,^2\text{He})$ reaction at $E_d = 170$ MeV at KVI [5] is a singlet state mixed with the unpolarized (random contamination) state with γ ($0 \leq \gamma \leq 1$) controlling the degree of mixing (the details of the experimental setup and analysis method of the $(d,^2\text{He})$ reaction

²In this case the directions $\hat{b}$ and $\hat{b}'$ of the analyser setting are equal to $\cos(\theta)\hat{c}_{\max} \pm \sin(\theta)\hat{c}'_{\max}$ and the direction $\hat{a}$, $\hat{a}'$ are equal to $\frac{D\hat{c}_{\max}}{\|D\hat{c}_{\max}\|}$, $\frac{D\hat{c}'_{\max}}{\|D\hat{c}'_{\max}\|}$, respectively. $\hat{c}_{\max}$ and $\hat{c}'_{\max}$ are two unit (not unique) and mutually orthogonal vectors in $\mathcal{R}^3$ that maximize the function $\|D\hat{c}\|^2 + \|D\hat{c}'\|^2$ (see Ref. [4] for more detail).




Figure 1: Plot of $\langle \mathcal{B} \rangle_{Werner}^{max}$ (dashed line) and τ (solid line) versus the purity γ . The dotted line is the Bell limit, the circle is the KVI limit for $\gamma \sim 0.9$.

were described in detail in Ref. [5]). Given all that, we can write the density matrix of such state as

$$\rho_W = (1 - \gamma) \frac{I}{4} + \gamma |\Psi^-\rangle \langle \Psi^-| \quad (10)$$

which interpolates between the unpolarized state $I/4$ and singlet state $|\Psi^-\rangle = (|+-\rangle - |-+\rangle)/\sqrt{2}$. This class of states is called Werner states [7]. The purity of Werner states is a monotonic function of γ . Thus, in this paper we use γ as our measure of purity. Also, for Werner state it is easy to prove using the condition noted above that

$$\langle \mathcal{B} \rangle_{Werner}^{max} = \gamma \langle \mathcal{B} \rangle_{pure}^{max}. \quad (11)$$

Note that, a violation of the modified Bell-inequality does not exclude an explanation with a hidden variable theory. In Fig. 1 we plot $\langle \mathcal{B} \rangle_{Werner}^{max}$ and the tangle τ versus the purity γ . As we can see in this figure the Werner state does not violate the Bell inequality if its purity γ is less than $1/\sqrt{2} \sim 70\%$. However, the entanglement is still non-zero in the Werner state until $\gamma > 1/3 \sim 33\%$. Therefore, some quantum correlation cannot be seen only

Table 1: Experimental data and quantum theory predictions for a pure singlet states (case 1) and mixed Werner states (case 2) for several violating cases of the CHSH-Bell inequality according to the definition given in Eq. 1.

CHSH-Bell Inequality	QM case 1	QM case 2	Exp. Data
$E(50^\circ, 0^\circ, 25^\circ, 75^\circ)$	2.46	2.21	0.67 ± 2.30
$E(60^\circ, 0^\circ, 30^\circ, 90^\circ)$	2.60	2.34	1.21 ± 2.42
$E(70^\circ, 0^\circ, 35^\circ, 105^\circ)$	2.72	2.45	1.54 ± 2.76
$E(80^\circ, 0^\circ, 40^\circ, 120^\circ)$	2.80	2.52	2.11 ± 2.86
$E(90^\circ, 0^\circ, 45^\circ, 135^\circ)$	2.83	2.55	2.23 ± 2.48
$E(100^\circ, 0^\circ, 50^\circ, 150^\circ)$	2.79	2.51	2.39 ± 2.87
$E(110^\circ, 0^\circ, 55^\circ, 165^\circ)$	2.69	2.34	2.58 ± 2.91
$E(120^\circ, 0^\circ, 60^\circ, 180^\circ)$	2.50	2.25	2.75 ± 2.95
χ^2	1.26	0.85	

by measuring the violation of the Bell-type inequality because some of them (Werner states) are entangled but still do not violate Bell inequality. Note that there is a possible experimental measurement of the entanglement based on the entanglement witness [8] that we think to implement in the future experiment.

In Tab. 1 we compare the quantum theory predictions assuming a pure singlet state (case 1) and mixed Werner states (case 2) for the spin of the two detected protons for several violating cases of the CHSH-Bell inequality. The value of $\chi^2 = \sum_i (\frac{R_{th}^i - R_{exp}^i}{\Delta R_{exp}^i})^2$ is given in the bottom of the table for both cases. We have found that $\chi_{Werner}^2 < \chi_{Singlet}^2$ as expected. However, we cannot judge this result as evidence of the mixing of the singlet with the unpolarized state because the experimental data suffer from large errors.

4 Conclusion

In this paper we have discussed the effect of the unpolarized state in the spin correlations measurement of the $^1S_0$ two proton state produced in $^{12}\text{C}(d, ^2\text{He})$ reaction at KVI. We have shown that even a small coupling (less than 10%) of the pure singlet state with the unpolarized state changes dramatically the Bell-violation value. After introducing the contribution of the unpolarized state we have found a better χ^2 . The experimental results are suffering from a large statistical error and therefore not conclusive for testing Bell's inequality, but with a modified experimental setup, measurements with significantly improved precision will become feasible.

Acknowledgments

This work was performed as part of the research program of the *Stichting voor Fundamenteel Onderzoek der Materie (FOM)* with financial support from the *Nederlandse Organisatie voor Wetenschappelijk Onderzoek*.

References

- [1] S. Hamieh and H. Zaraket, *J. Phys. A: Math. Gen.* **36**, L387-L391 (2003).
- [2] J.S. Bell, *Physics* **1**, 195 (1964); *Rev. Mod. Phys.* **38**, 447 (1966).
- [3] J. Clauser, M. Horne, A. Shimony, and R. Holt, *Phys. Rev. Lett.* **23**, 880 (1969).
- [4] R. Horodecki, P. Horodecki, and M. Horodecki, *Phys. Lett. A* **200**, 340 (1995).
- [5] C. Polachic, C. Rangacharyulu, A. van den Berg, M. Harakeh, M. de Huu, H. Wortche, J. Heyse, C. Baumer, D. Frekers, and J. Brooke, *quant-ph/0303136*, submitted to *Phys. Lett. A*.
- [6] C. H. Bennett, D. P. DiVincenzo, J. A. Smolin and W. K. Wootters, *Phys. Rev. A* **54**, 3824 (1996).
- [7] R.F. Werner, *Phys. Rev. A* **40**, 4277 (1989).
- [8] O. Guhne, P. Hyllus, D. Brub, E. Ekert, M. Lewenstein, C. Macchiavello, and A. Sampera, *Phys. Rev. A* **66**, 062305 (2002).